

The Commonwealth of Massachusetts

PRESENTED BY:

David Henry Argosky LeBoeuf and Mindy Domb

To the Honorable Senate and House of Representatives of the Commonwealth of Massachusetts in General Court assembled:

The undersigned legislators and/or citizens respectfully petition for the adoption of the accompanying bill:

An Act relative to reducing the risk of unintended fatal opioid overdose.

PETITION OF:

NAME:	DISTRICT/ADDRESS:	DATE ADDED:
David Henry Argosky LeBoeuf	17th Worcester	2/19/2021
Mindy Domb	3rd Hampshire	2/19/2021
Christina A. Minicucci	14th Essex	2/26/2021
Vanna Howard	17th Middlesex	2/26/2021
Nika C. Elugardo	15th Suffolk	2/26/2021
Elizabeth A. Malia	11th Suffolk	3/15/2021
David Allen Robertson	19th Middlesex	3/16/2021

HOUSE DOCKET, NO. 3723 FILED ON: 2/19/2021

By Representatives LeBoeuf of Worcester and Domb of Amherst, a petition (accompanied by bill, House, No. 2110) of David Henry Argosky LeBoeuf, Mindy Domb and others relative to reducing the risks of unintended fatal opioid overdoses. Mental Health, Substance Use and Recovery.

The Commonwealth of Massachusetts

In the One Hundred and Ninety-Second General Court (2021-2022)

An Act relative to reducing the risk of unintended fatal opioid overdose.

Be it enacted by the Senate and House of Representatives in General Court assembled, and by the authority of the same, as follows:

1	Chapter 111 of the General Laws is hereby amended by adding the following section:-
2	Section 238. Prior to issuing a prescription of an opioid which is a controlled substance as set
3	forth in section 31 of chapter 94C, a practitioner shall discuss with the patient, or the patient's
4	parent or guardian if the patient is under 18 years of age and is not an emancipated minor, the
5	risks for unintentional opioid overdose associated with the medication, including but not limited
6	to: (i) taking more opioids than prescribed; (ii) the dangers of mixing opioids with alcohol,
7	benzodiazepines such as Klonopin, Valium, and Xanax, other central nervous system
8	depressants, and other forms of poly-drug use; (iii) alternative treatments that may be available.
9	The physician will also discuss the risk of opioid dependence as a result of prescription use.
10	To avoid unintentional fatal overdose, the practitioner shall reconcile all patient
11	medications prior to prescribing opioids. The practitioner will require the patient to sign an
12	acknowledgement that the patient, or the patient's parent or guardian, as applicable, has (i)

- 13 discussed with the practitioner the risks of developing a physical or psychological dependence on
- 14 opioids, and (ii) understands the ways in which their alcohol use or other prescriptions may
- 15 increase the risk for fatal opioid overdose.